

## ALGEBRAS IN $MC_n(k)$ WITH $\dim(m_R^2) = 1$

YOUNGKWON SONG\*

ABSTRACT. We introduce a method to construct some algebras  $R \in MC_n(k)$  with  $\dim(R) = n$  and  $\dim(m_R^2) = 1$  for each  $n \geq 3$ .

### 1. Introduction

Throughout this paper,  $(R, m_R, k)$  is a local maximal commutative subalgebra of matrix algebra  $M_n(k)$  of size  $n \times n$  with nilpotent maximal ideal  $m_R$  and residue class field  $k$ . The set of all local maximal commutative subalgebras  $(R, m_R, k)$  of  $M_n(k)$  will be denoted by  $MC_n(k)$ . We assume the algebra  $R \in MC_n(k)$  contains the multiplicative identity. The socle of the algebra  $R$  is denoted by  $\text{soc}(R)$ . Furthermore,  $I_t$  is the identity matrix of size  $t \times t$  and  $O_{t \times s}$  is the zero matrix of size  $t \times s$ .

The next theorems are known as the Kravchuk's theorem.

**THEOREM 1.1.** ([5] Kravchuk's first theorem) *Let  $(R, m_R, k)$  be an algebra in  $MC_n(k)$ . Then, the matrix  $r \in m_R$  can be assumed to be of the following form :*

$$r = \begin{pmatrix} O_{\ell \times \ell} & O & O \\ A(r) & B(r) & O \\ C(r) & D(r) & O_{q \times q} \end{pmatrix},$$

where  $B(r) \in M_p(k)$ ,  $n = \ell + p + q$ ,  $\ell \neq 0, p \neq 0, q \neq 0$ . Moreover,  $\text{soc}(R)$  consists of all matrices of the form :

$$r = \begin{pmatrix} O_{(n-q) \times \ell} & O_{(n-q) \times (n-\ell)} \\ C(r) & O_{q \times (n-\ell)} \end{pmatrix},$$

where  $C(r) \in M_{q \times \ell}(k)$ .

---

Received June 22, 2011; Accepted August 13, 2011.

2010 Mathematics Subject Classification: Primary 15A27, 15A33.

Key words and phrases: maximal commutative subalgebra, socle.

The present research has been conducted by the Research Grant of Kwangwoon University in 2010.

THEOREM 1.2. ([5] Lemma 6) *Let  $(R, m_R, k)$  be an algebra in  $MC_n(k)$ . Suppose the matrices  $r_i \in m_R$  which are of the form*

$$r_i = \begin{pmatrix} O_{\ell \times \ell} & O & O \\ A(r_i) & B(r_i) & O \\ C(r_i) & D(r_i) & O_{q \times q} \end{pmatrix}, \quad i = 1, 2, \dots, t$$

*constitute a basis for  $m_R$  where  $B(r_i) \in M_p(k)$ . Then, the rank of the following  $p \times \ell t$  matrix  $H$  is  $p$  :*

$$H = \begin{pmatrix} A(r_1) & A(r_2) & \cdots & A(r_t) \end{pmatrix}.$$

THEOREM 1.3. ([1] Theorem 4 ) *Let  $(R, m_R, k)$  be a commutative algebra. Then,  $R$  is a  $C_1$ -construction if and only if there is an ideal  $N$  of  $R$  satisfying the following conditions :*

- (1)  $Ann_R(N) = N$
- (2) *The exact sequence  $0 \rightarrow N \rightarrow R \rightarrow R/N \rightarrow 0$  splits as  $k$ -algebras, where  $Ann_R(N)$  is the annihilator of  $N$ .*

Also, theorem 1.4 is an equivalent condition for a algebra  $R$  to be an algebra of the  $C_2$ -construction. The proof can be found in [3].

THEOREM 1.4. ([3] Lemma 2.8) *Let  $(R, m_R, k)$  be a finite dimensional commutative algebra. Then,  $R$  is a  $C_2$ -construction if and only if  $R$  contains a subalgebra  $(B, m_B, k)$  and an element  $x \in m_R$  satisfying the following conditions :*

- (1)  $x^\nu \neq 0 \in soc(B)$  for some positive integer  $\nu > 1$
- (2)  $m_B x = \{0\}$
- (3)  $dim_k(R) = dim_k(B) + (\nu - 1)$

The following theorem 1.5 is an equivalent condition to be a  $C_2^t$ -construction that can be found in [9].

THEOREM 1.5. ([9] Theorem 3.1) *Let  $(R, m_R, k)$  be a finite dimensional commutative algebra and let  $t$  be a positive integer. Then,  $R$  is a  $C_2^t$ -construction if and only if there exist a subalgebra  $(B, m_B, k)$  of  $R$  and elements  $x_i \in m_R$ ,  $i = 1, 2, \dots, t$  satisfying the following properties :*

- (1)  $x_i^2 = x_j^2 \in soc(B) - \{0\}$  for all  $1 \leq i, j \leq t$
- (2)  $x_i x_j = 0$  for all  $1 \leq i \neq j \leq t$
- (3)  $m_B x_i = \{0\}$  for all  $1 \leq i \leq t$
- (4)  $dim_k(R) = dim_k(B) + t$

## 2. Algebras in $MC_n(k)$ with $\dim(R) = n$ and $\dim(m_R^2) = 1$

In this section, for each positive integer  $n \geq 3$ , we will introduce a construction to produce some algebras  $R \in MC_n(k)$  with  $\dim(R) = n$  and  $\dim(m_R^2) = 1$ .

From now on, we will consider the case of  $\ell = 1$  in theorem 1.1. That is, we assume the matrices  $r_i$  in  $m_R$  are of the form

$$r_i = \begin{pmatrix} O_{1 \times 1} & O & O \\ A(r_i) & B(r_i) & O \\ C(r_i) & D(r_i) & O_{q \times q} \end{pmatrix}, \quad i = 1, 2, \dots, p + q$$

constitute a basis for  $m_R$ .

Now, we define  $\theta$ -relation as follows :

DEFINITION 2.1. Let  $\theta$  be a subset of a set  $\{1, 2, \dots, q\}$ . We say the pair  $(A(r_i), D(r_i))$  are in  $\theta$ -relation if for each  $A(r_i) \in M_{p \times 1}(k)$ , the matrix  $D(r_i) \in M_{q \times p}(k)$  are defined as follows :

$$D(r_i) = \begin{pmatrix} D_1(r_i) \\ D_2(r_i) \\ \vdots \\ D_q(r_i) \end{pmatrix}, \quad D_j(r_i) = \begin{cases} A(r_i)^T, & \text{if } j \in \theta, \\ O_{1 \times p}, & \text{otherwise} \end{cases}$$

Here,  $A(r_i)^T$  is the transpose of  $A(r_i)$ .

Now, we can construct algebras  $R \in MC_n(k)$  with  $\dim(R) = n$  and  $\dim(m_R^2) = 1$  for each  $n \geq 3$  as the following theorem.

THEOREM 2.2. Let  $R$  be a subalgebra of  $M_n(k)$  and let  $m_R$  have a basis consisting of following form of matrices :

$$r_i = \begin{pmatrix} O_{1 \times 1} & O & O \\ A(r_i) & O_{p \times p} & O \\ C(r_i) & D(r_i) & O_{q \times q} \end{pmatrix}, \quad i = 1, 2, \dots, p + q$$

, where  $A(r_i) \in M_{p \times 1}(k)$ ,  $C(r_i) \in M_{q \times 1}(k)$ ,  $D(r_i) \in M_{q \times p}(k)$ , and  $n = p + q + 1$ . If the pairs  $(A(r_i), D(r_i))$  are in  $\theta$ -relation for all  $i = 1, 2, \dots, p + q$ , then  $R$  is an algebra in  $MC_n(k)$ .

*Proof.* Note that by theorem 1.1 and theorem 1.2, we may assume  $A(r_i)$ ,  $A(r_j)$  and  $C(r_i)$ ,  $C(r_j)$  are of the following form for  $i = 1, 2, \dots, p$ ,  $j = p + 1, p + 2, \dots, p + q$ :

$$A(r_i) = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix} \leftarrow i^{\text{th}} \text{ row}, \quad C(r_j) = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix} \leftarrow j^{\text{th}} \text{ row}$$

$$A(r_j) = O_{p \times 1}, \quad C(r_i) = O_{q \times 1}$$

Now let

$$S = \begin{pmatrix} S_{11} & S_{12} & S_{13} \\ S_{21} & S_{22} & S_{23} \\ S_{31} & S_{32} & S_{33} \end{pmatrix} \in M_n(k)$$

Here,  $S_{11} \in k$ ,  $S_{22} \in M_p(k)$ ,  $S_{33} \in M_q(k)$ .

Then, we have the following equations from the equation  $r_i S = S r_i$  for all  $i$  :

- (1)  $S_{12}A(r_i) + S_{13}C(r_i) = 0$
- (2)  $S_{13}D(r_i) = 0$
- (3)  $S_{22}A(r_i) + S_{23}C(r_i) = A(r_i)S_{11}$
- (4)  $S_{23}D(r_i) = A(r_i)S_{12}$
- (5)  $A(r_i)S_{13} = O_{p \times q}$
- (6)  $S_{32}A(r_i) + S_{33}C(r_i) = C(r_i)S_{11} + D(r_i)S_{21}$
- (7)  $S_{33}D(r_i) = C(r_i)S_{12} + D(r_i)S_{22}$
- (8)  $C(r_i)S_{13} + D(r_i)S_{23} = O_{q \times q}$

From the equation (1) and (3),  $S_{12} = O_{1 \times p}$ ,  $S_{13} = O_{1 \times q}$ ,  $S_{23} = O_{p \times p}$ . Thus, we have the following equations :

- (3-1)  $S_{22}A(r_i) = A(r_i)S_{11}$
- (6-1)  $S_{33}C(r_i) = C(r_i)S_{11}$
- (6-2)  $S_{32}A(r_i) = D(r_i)S_{21}$

From the equation (3-1),

$$S_{22}A(r_i) = \text{Col}_i(S_{22}) = \begin{pmatrix} 0 \\ \vdots \\ s_{11} \\ \vdots \\ 0 \end{pmatrix} \leftarrow i^{\text{th}} = A(r_i)S_{11},$$

where  $S_{11} = (s_{11})_{1 \times 1}$  and  $\text{Col}_i(S_{22})$  is the  $i$ -th column of  $S_{22}$ . Thus,  $S_{22} = s_{11}I_p$ .

From the equation (6-1),

$$S_{33}C(r_i) = Col_i(S_{33}) = \begin{pmatrix} 0 \\ \vdots \\ s_{11} \\ \vdots \\ 0 \end{pmatrix} \leftarrow i^{th} = C(r_i)S_{11}$$

Thus,  $S_{33} = s_{11}I_q$ .

From the equation (6-2), we have  $Col_i(S_{32}) = S_{32}A(r_i) = D(r_i)S_{21}$ .

If we let  $S_{21} = (d_1, d_2, \dots, d_p)^T$  for some  $d_i \in k$ ,  $i = 1, 2, \dots, p$ , then

$$D(r_i)S_{21} = \begin{pmatrix} D_1(r_i) \\ D_2(r_i) \\ \vdots \\ D_q(r_i) \end{pmatrix} \begin{pmatrix} d_1 \\ d_2 \\ \vdots \\ d_p \end{pmatrix} = \begin{pmatrix} d_{1i} \\ d_{2i} \\ \vdots \\ d_{qi} \end{pmatrix},$$

where  $d_{ti} = \begin{cases} d_i, & \text{if } t \in \theta, \\ 0, & \text{otherwise} \end{cases}$

Thus,

$$Col_i(S_{32}) = \begin{pmatrix} d_{1i} \\ d_{2i} \\ \vdots \\ d_{qi} \end{pmatrix}$$

and so the matrix  $S$  is of the form

$$S = \begin{pmatrix} a & O & O \\ S_{21} & aI_p & O \\ S_{31} & S_{32} & aI_q \end{pmatrix}.$$

for some  $a \in k$ , where

$$S_{21} = \begin{pmatrix} d_1 \\ d_2 \\ \vdots \\ d_p \end{pmatrix}, \quad S_{32} = \begin{pmatrix} S_1 \\ S_2 \\ \vdots \\ S_q \end{pmatrix},$$

where

$$S_j = \begin{cases} S_{21}^T, & \text{if } j \in \theta, \\ O_{1 \times p}, & \text{otherwise} \end{cases}$$

Here,  $S_{21}^T$  is the transpose of  $S_{21}$ . This implies  $S \in R$  and therefore, we can conclude that  $R \in MC_n(k)$ .  $\square$

EXAMPLE 2.3. Let  $R$  be a subalgebra of  $M_6(k)$  defined as following:

$$R = \left\{ \begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 \\ a_1 & a & 0 & 0 & 0 & 0 \\ a_2 & 0 & a & 0 & 0 & 0 \\ a_3 & a_1 & a_2 & a & 0 & 0 \\ a_4 & 0 & 0 & 0 & a & 0 \\ a_5 & a_1 & a_2 & 0 & 0 & a \end{pmatrix} \mid a, a_1, a_2, a_3, a_4, a_5, \alpha \in k \right\}$$

Then, we may consider as  $p = 2, q = 3, \theta = \{1, 3\}$ ,

$$\begin{aligned} A(r_1) &= \begin{pmatrix} 1 \\ 0 \end{pmatrix}, & A(r_2) &= \begin{pmatrix} 0 \\ 1 \end{pmatrix}, & A(r_3) &= A(r_4) = A(r_5) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \\ D(r_1) &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} A(r_1)^T \\ O \\ A(r_1)^T \end{pmatrix}, & D(r_2) &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} A(r_2)^T \\ O \\ A(r_2)^T \end{pmatrix}, \\ D(r_3) &= D(r_4) = D(r_5) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \\ C(r_1) &= C(r_2) = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, & C(r_3) &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \\ C(r_4) &= C(r_2) = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, & C(r_5) &= \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \end{aligned}$$

Thus, by the theorem 2.2,  $R$  should be a subalgebra in  $MC_6(k)$  with  $\dim(R) = 6$  and  $\dim(m_R^2) = 1$ .

The following lemma 2.4 and theorem 2.5 provides some properties of an algebra  $R \in MC_n(k)$  in theorem 2.2 and the proof can be obtained by straightforward calculations. The  $(i, j)^{th}$  matrix unit will be denoted by  $E_{ij}$ .

LEMMA 2.4. Suppose  $R \in MC_n(k)$  is an algebra as in theorem 2.2. Then,  $m_R^2 = (E_{1+p+\ell_1, 1} + \dots + E_{1+p+\ell_\mu, 1})$ , where  $\ell_1, \dots, \ell_\mu \in \theta$  with  $\ell_1 < \dots < \ell_\mu$ .

THEOREM 2.5. Suppose  $R \in MC_n(k)$  is an algebra as in theorem 2.2. Then, the following properties hold :

- (1)  $\dim(R) = n$

- (2)  $\dim(m_R^2) = 1$
- (3)  $m_R^2 \subseteq \text{soc}(R)$
- (4)  $\dim(\text{soc}(R)) = q$
- (5)  $\dim(\text{soc}(R)/m_R^2) = q - 1$
- (6)  $i(m_R) = 3$ , where  $i(m_R)$  is the index of the nilpotency of  $m_R$ .

### 3. Relation with $C_i$ -constructions

In this section, we want to prove if the construction in section 2 imply the  $C_2$ -construction and the  $C_2^t$ -construction but not the  $C_1$ -construction.

**THEOREM 3.1.** *Suppose  $R \in MC_n(k)$  is an algebra in theorem 2.2. Then,  $R$  is not a  $C_1$ -construction.*

*Proof.* Suppose  $R$  is a  $C_1$ -construction. Then,  $R$  should contain an ideal  $N$  satisfying  $\text{Ann}_R(N) = N$ . Let  $r \in \text{Ann}(N)$ . Then  $r = a_1r_1 + a_2r_2 + \dots + a_pr_p + bs$  for some  $a_i, b \in k$ ,  $i = 1, 2, \dots, p$  and  $s \in \text{soc}(R)$ . Note that  $r^2 = (a_1^2 + a_2^2 + \dots + a_p^2)s_1$  for some  $s_1 \neq O_{n \times n} \in \text{soc}(R)$ . Since  $r \in \text{Ann}_R(N)$ ,  $r^2 = O_{n \times n}$  and so  $r^2 = (a_1^2 + a_2^2 + \dots + a_p^2)s_1 = O_{n \times n}$  which implies  $a_i = 0$  for all  $i = 1, 2, \dots, p$ . Thus,  $r = bs \in \text{soc}(R)$  and so  $\text{Ann}_R(N) \subseteq \text{soc}(R)$ . Since  $\text{soc}(R) \subseteq \text{Ann}_R(N)$ , we have  $N = \text{Ann}_R(N) = \text{soc}(R)$ . But then  $N = \text{Ann}_R(\text{soc}(R)) = m_R$  and  $m_R^2 = N^2 = \{O_{n \times n}\}$  which is impossible since  $\dim(m_R^2) = 1$ . Therefore, there doesn't exist an ideal  $N$  satisfying  $\text{Ann}_R(N) = N$  and we can conclude that  $R$  is not a  $C_1$ -construction.  $\square$

**THEOREM 3.2.** *Suppose  $R \in MC_n(k)$  is an algebra in theorem 2.2. Then,  $R$  is a  $C_2$ -construction.*

*Proof.* Let  $B = k[r_2, \dots, r_p] \oplus \text{soc}(R)$ . Then  $B$  is a subalgebra of  $R$  and for the element  $x = r_1$ , the following properties holds :

- (1)  $x^2 \neq O_{n \times n} \in \text{soc}(B)$
- (2)  $m_Bx = \{O_{n \times n}\}$
- (3)  $\dim(R) = \dim(B) + 1$ .

Thus, the algebra  $R$  satisfies the conditions in theorem 1.4 and so  $R$  is a  $C_2$ -construction.  $\square$

**THEOREM 3.3.** *Suppose  $R \in MC_n(k)$  is an algebra in theorem 2.2. Then,  $R$  is a  $C_2^t$ -construction.*

*Proof.* Let  $B = k[\text{soc}(R)]$  and let  $x_i = r_i$  for all  $i = 1, 2, \dots, p$ . Then  $B$  and  $x_i$  satisfies the following conditions :

- (1)  $x_i^2 = x_j^2 \in \text{soc}(B) - \{O_{n \times n}\}$  for all  $1 \leq i, j \leq p$
- (2)  $x_i x_j = O_{n \times n}$  for all  $1 \leq i \neq j \leq p$
- (3)  $m_B x_i = \{O_{n \times n}\}$  for all  $1 \leq i \leq p$
- (4)  $\dim_k(R) = \dim_k(B) + p$

Thus, by the theorem 1.5,  $R$  is a  $C_2^t$ -construction for  $t = p$ .  $\square$

### References

- [1] W. C. Brown and F. W. Call, *Maximal Commutative Subalgebras of  $n \times n$  Matrices*, Communications in Algebra **59** (1993), no. 12, 4439–4460.
- [2] W. C. Brown, *Two Constructions of Maximal Commutative Subalgebras of  $n \times n$  Matrices*, Communications in Algebra **22** (1994), no. 10, 4051–4066.
- [3] ———, *Constructing Maximal Commutative Subalgebras of Matrix Rings in Small Dimensions*, Communications in Algebra **25** (1997), no. 12, 3923–3946.
- [4] R. C. Courter, *The Dimension of Maximal Commutative Subalgebras of  $K_n$* , Duke Mathematical Journal **32** (1965), 225–232.
- [5] D. A. Suprunenko and R. I. Tyshkevich, *Commutative Matrices*, Academic Press, 1968.
- [6] Youngkwon Song, *On the Maximal Commutative Subalgebras of 14 by 14 Matrices*, Communications in Algebra **25** (1997), no. 12, 3823–3840.
- [7] ———, *Maximal Commutative Subalgebras of Matrix Algebras*, Communications in Algebra **27** (1999), no. 4, 1649–1663.
- [8] ———, *Notes on the Constructions of Maximal Commutative Subalgebra of  $M_n(k)$* , Communications in Algebra **29** (2001), no. 10, 4333–4339.
- [9] ———, *On  $C_2^t$ -Construction*, FJMS **33** (2009), no. 1, 33–40.

\*

Department of Mathematics  
 Kwangwoon University  
 Seoul 139–701, Republic of Korea  
*E-mail:* yksong@kw.ac.kr